



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: A61K 9/00, 31/575	A1	(11) International Publication Number: WO 98/13023 (43) International Publication Date: 2 April 1998 (02.04.98)
(21) International Application Number: PCT/FI97/00585 (22) International Filing Date: 26 September 1997 (26.09.97) (30) Priority Data: 963904 27 September 1996 (27.09.96) FI (71) Applicant (for all designated States except US): SUOMEN SOKERI OY [FI/FI]; Kyllikinportti 2, FIN-00240 Helsinki (FI). (72) Inventors; and (75) Inventors/Applicants (for US only): TIAINEN, Ilkka [FI/FI]; Pohjoiskaari 19 F 21, FIN-00200 Helsinki (FI). NURMI, Juha [FI/FI]; Vasarasepantie 4 D 6, FIN-10330 Pinjainen (FI). KLASI, Leena [FI/FI]; Intalantie 17, FIN-31600 Jokioinen (FI). (74) Agent: KOLSTER OY AB; Iso Roobertinkatu 23, P.O. Box 148, FIN-00121 Helsinki (FI).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>In English translation (filed in Finnish).</i>
(54) Title: A PRODUCT, A METHOD FOR ITS PRODUCTION, AND ITS USE		
(57) Abstract		
<p>The invention relates to a product containing a microcrystalline plant sterol, a method for producing the product by pulverization, and use of the product for the manufacture of edible products. The invention also relates to products manufactured using the microcrystalline plant-sterol-containing product. Preferably, the product is a spread based on a combination of a sweetening agent, a microcrystalline plant sterol and suitable berry or fruit.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

A product, a method for its production, and its use

Field of the invention

5 The invention relates to a product containing a microcrystalline plant sterol, to a method for producing the product through pulverization, and to the use of the product for producing edible products. The invention also relates to products produced by using this product containing a microcrystalline plant sterol. A preferred
10 product is a spread based on a combination of a sweetening agent, a microcrystalline plant sterol and a suitable berry or fruit.

Background

15 The cholesterol-lowering effect of plant sterols has been known since the 1950's, and the literature of the field therefore includes several reports concerning the positive effects of plant sterols on the cholesterol level. New reports include for example the research by Heinemann et al., Eur. J. Clin. Pharmacol. 40 (Suppl. 1),
20 1991, 59-63, about the effect of sitostanol and sitosterol on the absorption of cholesterol in the intestine by using the intestinal perfusion technique. Sitostanol was found to reduce the cholesterol absorption by as much as 85% and the sitosterol absorption by 50%.

25 Nevertheless, the use of plant sterols has not been very extensive, due to for example their poor availability and therefore high price. One of the greatest problems has been the poor solubility of plant sterols, since they are not soluble in water and are also poorly
30 soluble in fats. There have been efforts to solve the solubility problem by forming fat-soluble forms of plant sterols, for example fatty acid esters, and by using them in connection with fats.

35 An example of a method for preparing a fatty acid ester is the method disclosed in EP 195,311 (Yoshikawa Oil

& Fat Co., Ltd.) wherein a plant sterol and a branched aliphatic primary or secondary alcohol having 14 to 32 carbon atoms is reacted with a fatty acid or fatty acid ester in the presence of lipase or cholesterol esterase in an aqueous solution or in a water-containing organic solvent. DE 2,035,069 (Procter & Gamble Co.) discloses the preparation of carboxylic acid esters of plant sterols with an acylation reaction by using perchloric acid as a catalyst. The obtained plant sterol ester is then added to a cooking or salad oil to produce an oil that reduces the cholesterol level.

WO 92/19640 (Raision Margariini Oy) discloses a β -sitostanol fatty acid ester that is prepared by esterifying a β -sitostanol mixture with a fatty acid ester or ester mixture in the presence of an interesterification catalyst. The obtained fatty acid ester is suggested to be used as a part of fats or oils in fat-containing products; a preferred embodiment is the addition of the ester to edible fats. The reference also discloses by way of comparison the effect of β -sitostanol that has been emulsified into rape-seed oil on serum cholesterol levels. The result is not statistically significant and it is considered marginal as compared with the effect of the β -sitostanol ester described in the reference.

EP 289,636 (Asahi Denka Kogyo Kabushiki Kaisha) relates to an emulsified or solubilized sterol composition wherein the sterols are emulsified or solubilized in an aqueous solution of polyhydroxy compounds containing sucrose and/or polyglycerol fatty acid esters or in liquid polyhydroxy compounds containing these fatty acid esters. The reference discloses that no solvent, such as fat or oil, is used for dissolving the sterols. However, the composition contains as an essential ingredient sucrose and/or polyglycerol fatty acid esters that are used in an amount of 0.5 to 10 parts by weight per part by weight of

the sterol. The product is stated to be stable and useful for example in foodstuffs, cosmetics, drugs and agricultural chemicals. These final products are not described in greater detail.

5 US 5,244,887 (Straub) discloses food additives containing plant stanols. The stanols include sitostanol, clionastanol, 22,23-dihydrobrassicastanol, campestanol, and mixtures thereof. In order to prepare the food
10 additive, a solution, suspension or emulsion of stanols is prepared by mixing the stanol with a solubilizing agent, an antioxidant and a dispersing agent. A preferred product is a composition containing about 74.8% vegetable oil, about 1.2% tocopherol and about 25% stanols. The additive
15 is added to foodstuffs containing cholesterol, for example to meat, eggs and dairy products, and the purpose is to prevent the absorption of cholesterol from these foodstuffs.

 The background art thus describes efforts to improve the solubility and therefore the range of use of
20 plant sterols by forming fat-soluble derivatives from the sterols; the plant sterols and the aforementioned derivatives are dissolved or emulsified in a fat or a fat component and they are used as additives in fat-containing foodstuffs to prevent the absorption of cholesterol from
25 these foodstuffs.

Description of the invention

 The object of the present invention was to provide such a form of a plant sterol that is useful as
30 such and that can be produced in a simple and economic manner without harmful additives.

 According to the present invention, these aims are achieved by using a microcrystalline plant sterol instead of a crystalline plant sterol.

 The invention therefore relates to a product
35 containing a crystalline plant sterol, the product being

characterized in that the plant sterol is in a microcrystalline form.

The volumetric mean particle size of the product according to the invention containing a microcrystalline plant sterol is less than 35 μm , preferably less than 30 μm and more preferably less than 25 μm , for example 20 to 23 μm . The volumetric mean particle size is most preferably a great deal lower than these values, for example about 4 to 15 μm , in particular 5 to 10 μm . When the volumetric mean particle size is about 10 to 20 μm , preferably at most 15% of the particles have a size exceeding 30 μm .

The particle size and volume distributions of a plant sterol pulverized into a microcrystalline form according to the invention and a crystalline plant sterol that has not been pulverized and that is used as reference material are shown in Figures 1 and 2, respectively. The assays were carried out by the PAMAS method in an aqueous solution to which a small amount of tenside had been added. The difference between the microcrystalline plant sterol according to the invention (Example 3) and the untreated crystalline plant sterol is very clear. The size and volume distributions of the non-pulverized sample are very broad, including both very small particles and a great number of large particles that prevent the addition of a crystalline plant sterol as such to edible products, causing both technical problems and problems with taste and structure. The microcrystalline plant sterol according to the invention is in turn clearly more homogenous and contains substantially fewer large particles.

A plant sterol here refers to a phytosterol that reduces the cholesterol level. The plant sterol is preferably β -sitosterol, i.e. 24-ethyl-5-cholesten-3 β -ol, or especially its hardened form β -sitostanol, i.e. 24-ethyl-5 α -cholesten-3 β -ol. Other suitable plant sterols

include for example campestenol and clionastanol. Commercially available plant sterols are often mixtures of plant sterols that are also appropriate for use according to the present invention.

5 A microcrystalline plant sterol can be formed by pulverization according to the present invention.

The invention therefore also relates to a method for preparing a product containing a crystalline plant sterol, the method being characterized in that the
10 crystalline plant sterol is pulverized into a microcrystalline form.

The pulverization can be carried out both as dry grinding and wet grinding. The dry grinding can be performed for example with a counterjet pulverizer wherein
15 the particles that have been fluidized into a carrier gas, typically air, and that are to be pulverized are conducted to collide at a high speed. The pulverizer preferably comprises a classifier where the pulverized material is classified into fractions having different particle sizes.
20 Other grinding apparatuses based on kinetic energy can also be used for carrying out the dry grinding. The dry grinding can also be carried out in mills based on the use of friction, for example in a ball mill. The wet grinding is performed preferably in mills of the aforementioned
25 type based on the use of friction. The parameters used in the grinding are determined according to the structure of the grinding device used and the substance or mixture to be pulverized.

The pulverization can be carried out on a
30 crystalline plant sterol as such or on a mixture of a plant sterol and another component, such as a suitable admixture. According to the present invention, the pulverization is carried out preferably on a mixture of a plant sterol and a sweetening agent.

The sweetening agent can be for example a normal sugar, such as sucrose, glucose or fructose, an aqueous solution of sugar, a starch syrup, isoglucose, isomalt, a sugar alcohol, such as sorbitol, lactitol or xylitol. The sugar can also be replaced with a mixture of a builder and a sweetening agent. The builder can be polydextrose or inulin and the sweetening agent can be an intense sweetener, a carbohydrate sweetener or a mixture thereof. The amount of the sweetening agent can vary depending on the method used and the desired final product. In sweet spreads, a sweetening agent and a plant sterol are usually used in the ratio of about 0.5:9.5 to 9.5:0.5, preferably about 3:7 to 9:1, most preferably about 4:6 to 6:4 and particularly about 1:1. The spreads do not always contain a sweetening agent. The function of sugar as a builder can be replaced in both sweet and other spreads with other builders. According to the present invention, a microcrystalline plant sterol can also be included in fat-containing spreads, such as margarine, cheese spread, pastes etc., having a low sugar content or containing no sugar.

According to the present invention, a separate homogenization stage can be carried out in addition to pulverization. The homogenization stage can be carried out both before and after the pulverization. According to the present invention, also the homogenization stage is carried out preferably on a mixture of a plant sterol and another component.

In a preferred embodiment of the present invention, a non-homogenous mixture or suspension of a plant sterol and a liquid phase, such as water or an aqueous solution, is formed first. This is converted into a homogenous presuspension by mixing for example with a blender preferably at a high mixing speed and/or by using a great shear force. It is preferable for the homogeneity

of the presuspension that the dry substance content and viscosity of the liquid phase are rather high. Thus it is preferable to prepare the premixture in a sugar-containing syrup instead of pure water. Most preferably, a liquid
5 sugar or a sugar syrup having a dry substance content of at least 65% is used. The proportion of the plant sterol is preferably 10% or more, particularly as much as 50% based on the dry substance content. It has been found out in connection with this invention that when a syrup
10 containing a plant sterol and sugar is mixed effectively, after a treatment of 20 minutes in a domestic blender the plant sterol crystals were completely wet and no crystal agglomerates were formed even after a long storage period.

The suspension is subjected thereafter to a
15 pulverization treatment wherein the plant sterol crystals in the suspension are pulverized into microcrystals. For example ball mills are suitable for this purpose, as stated above.

In a second preferred embodiment of the present
20 invention, especially in large-scale production, pulverization is carried out first, preferably on a mixture of a plant sterol and an admixture, such as a sweetening agent. The pulverized mixture can also be homogenized or suspended.

25 According to the present invention, a mainly mechanical treatment is carried out; no esterification is performed and no other derivatives are produced either, and no solvents or other non-edible admixtures are required. The method according to the invention is much
30 simpler than chemical methods, and the product will not contain any harmful residues of solvents, catalysts or admixtures. It should also be noted that the derivatives that are made highly soluble according to prior art methods can release a very poorly soluble sterol to the
35 organism when they decompose and possibly cause risks of

precipitation. Such significant changes in solubility do not occur when a microcrystalline plant sterol is used according to the present invention.

5 The product according to the invention containing
a microcrystalline plant sterol has a cholesterol-lowering
effect. The product can be used as such. It can also be
used as an ingredient in cholesterol-lowering products,
such as functional foodstuffs, natural products and
pharmaceuticals. A microcrystalline plant sterol
10 homogenized into a fat or an aqueous solution is
applicable for use especially in the manufacture of other
edible cholesterol-lowering products; the degree of
processing of the sterol is suitable for the industry and
no processing problems occur during its use.

15 When a microcrystalline plant sterol was added
according to the present invention to edible products, the
plant sterol had no harmful effects on the structure,
taste or mouth-feel of the final product. Unpleasant taste
or aftersensations reported earlier in connection with
20 plant sterols were not found.

The invention thus relates further to the use of
a product containing a microcrystalline plant sterol as a
cholesterol-lowering agent and to its use for the
manufacture of edible products.

25 The present invention also provides edible
cholesterol-lowering products, which contain a micro-
crystalline plant sterol according to the invention in
addition to the other conventional ingredients.

30 A spread based on a combination of a sweetening
agent, a microcrystalline plant sterol and a suitable
berry or fruit is disclosed in the invention as a
preferred edible cholesterol-lowering product. It has been
proved in connection with the present invention that it is
possible to use in the product a microcrystalline plant
35 sterol as such without any technological problems. There

are no problems in the product that can be detected by sensory evaluation and that relate to taste or structure. The product can be used in the same manner as fats or marmalade for example on bread or other bakery products. 5 The product can replace the use of butter or margarine and it has several advantageous characteristics over them.

Due to the plant sterol, the product has a cholesterol-lowering effect. The use of the product also directly reduces the total intake of fat since no fat, but 10 a non-fat spread, is put on bread. At the same time, especially the use of hardened fats can be reduced. Thus the proportion of fats and salt of the total intake of energy decreases, which is recommendable for serum cholesterol level and blood pressure. The product makes 15 the diet lighter, since the spread contains less than half of the energy content of butter or margarine and about 25 to 30% less energy than light fats having a fat content of 40%. The use of the product also decreases the intake of salt, since it does not require the addition of any salt, 20 whereas fats usually contain about 0.5 to 1.5% of salt. Decreasing the intake of salt and especially sodium is highly recommendable for health, as is well known. Compared with fats, the product also has a clearly longer shelf life.

25 The sweetening component of the preferred spread according to the invention can be a conventional sugar, such as sucrose for example in the form of normal granulated sugar, glucose or fructose, or special sugars, such as preservation sugar, containing for example a 30 jellying agent and/or a preservative. If desired, the sweetening component can be an aqueous solution of sugar or several sugars, a mixture of several sugars, or it can partly consist of products that are obtained during the refinement of sugar or that are manufactured specifically, 35 such as liquid sugars, sugar syrups or mixtures thereof.

The sweetening component of the product or a part of it can also be, if desired, for example starch syrup, isoglucose, polyol, such as sorbitol or xylitol, polydextrose or an intense sweetener in addition to the other usual sweetening agents.

Suitable berries for the preferred spread according to the present invention include, for example, strawberry, blueberry, boysenberry, lingonberry, gooseberry, raspberry, blackberry and currants. Examples of suitable fruits include, for example, orange and other citrus fruits, apricot, apple, peach, plum and cherry. It is evident that the spread according to the invention can also contain different combinations of berries and/or fruits.

The plant sterol can be selected, as stated above, from several different plant sterols or plant sterol mixtures. Preferable sterols are β -sitosterol and especially β -sitostanol.

The product can be supplemented, if desired, with usual additives and other additional agents, such as jelling and thickening agents, preservatives, acidity-regulating agents, flavourings, aromatic agents etc.

The spread according to the present invention can therefore have for example the following composition:

Fruit and/or berry	20 - 60%
Sweetening agent	20 - 60%
Plant sterol	0.5 - 10%
Jellying agent	0 - 5 %
Preservatives	0 - 0.1%
Water	ad 100%

The product preferably comprises about 30 to 50%, preferably 40 to 48% of fruit and/or berry, and about 30 to 55%, most preferably about 40 to 50% of a sweetening

agent. Pectin can preferably be used as the jellying agent. The preservatives can be usual preservatives, such as potassium sorbate. The product can also be supplemented, if desired, with other usual additives, such as citric acid, as an acidity-regulating agent. By changing the ingredients of the product and the method of manufacture the spread can be prepared for example in the form of jam, marmalade or jelly. In a corresponding manner, with the microcrystalline plant sterol according to the present invention it is also possible to produce beverages, such as nectars.

The cholesterol-lowering effect of a plant sterol can be increased by using pectin in the products of the invention that contain a microcrystalline plant sterol, usually in an amount of about 0.1 to 2%, preferably 0.2 to 1%, most preferably 0.4 to 0.6%. In addition to intensifying the effect of the plant sterol, pectin also has a positive effect on the product structure. It is also noted that the berries and/or fruits contained in the preferred spread according to the invention also inherently contain pectins.

A product corresponding to the spread according to the invention, based on a combination of a sweetening agent, a plant sterol and a suitable berry or fruit, can also be prepared by replacing the berries or fruits with molasses, such as a sugar syrup containing trace elements, or a mixture of sugar syrup and starch syrup. The product can be supplemented with a thickening agent, if desired. The structure of the product according to the invention that contains a microcrystalline plant sterol and that is based on molasses corresponds to honey or peanut butter and it has all the advantages of the above-described spread based on berries or fruits.

It is also possible to use sugar and different sweetening products, such as special sugars containing

small amounts of a jellying agent and/or a preservative and intended for jam-making or for baking, to manufacture, according to the present invention, corresponding functional cholesterol-lowering sweetening products by adding a microcrystalline plant sterol thereto. The edible cholesterol-lowering products according to the invention also include different sugar-containing products and spice-sugar mixtures to which a microcrystalline plant sterol has been added according to the present invention.

Examples of other functional products according to the invention that contain a sweetening agent and a microcrystalline plant sterol include different sweets, such as marmalades, chocolate, sugar candies and chewing gums; nectars and other beverages; different desserts based on milk or fruit, such as puddings and creams; yoghurts, salad dressings, such as mayonnaise, etc.

The sugar content of the functional products according to the invention that contain a sweetening agent can be about 1 to 99%, in preferred products about 45 to 95%. The functional sweetener products according to the invention contain about 0.5 to 15%, preferably about 1 to 10% of a plant sterol. It should be understood, however, that these values, as all the other numerical values given in this specification, are only trend-setting values. According to the invention, it is therefore possible to use all amounts sufficient for achieving the desired effect and having no negative effect on the structure or taste of the final product.

The microcrystalline plant sterol according to the invention can also be prepared and used without an added sweetening agent. Therefore also all the other edible cholesterol-lowering products containing a microcrystalline plant sterol fall within the scope of the present invention. They include especially products that are based on fats and that contain a microcrystalline

plant sterol, for example vegetable oil and/or fats or animal fats containing a microcrystalline plant sterol, or mixtures thereof. The microcrystalline plant sterol mixes well with fat-containing foodstuffs, such as soft margarine and baking margarine or mayonnaise, both as such and as a mixture of a plant sterol and a sweetening agent. It is easy to prepare a homogenous mixture also at room temperature. A microcrystalline plant sterol can also be added to bakery products and/or to products used in their preparation both as such and as a mixture with for example a sweetening agent, fat, flour or some other suitable component.

One of the most important advantages of edible cholesterol-lowering products according to the invention is that the plant sterol is brought to the consumers in such a form that is easy to use.

The edible cholesterol-lowering products according to the invention are prepared in the same manner as the corresponding conventional products by adding a desired amount of a microcrystalline plant sterol, which has optionally been homogenized, at a suitable stage of the preparation.

The microcrystalline plant sterol according to the invention is also applicable for use in pharmaceutical products, as already stated. These products are also prepared by methods that are common in the field, adding a desired amount of a microcrystalline plant sterol, which has optionally been homogenized, at a suitable stage of the preparation.

The invention will be described in greater detail by means of the examples below. The examples are only provided to illustrate the invention and they should not be considered to restrict the scope of the invention.

Example 1

Preparation of a homogenized microcrystalline plant sterol

150 g of sitostanol (Kaukas Oy) was added as such to 2310 g of Neste 65 syrup (Porkkalan Sokeripuhdistamo Oy). The mixture was homogenized with a blender (Braun, type 4259) for 30 min, whereafter the suspension was pulverized for 17 h in a large porcelain ball mill.

Example 2

Preparation of a homogenized microcrystalline plant sterol

50 g of sitostanol was added as such to 650 g of Neste 77 syrup (Porkkalan Sokeripuhdistamo Oy). The mixture was homogenized and a presuspension was formed with a blender (Braun, type 4259) for 30 min, whereafter the suspension was pulverized for 17 h in a large porcelain ball mill.

Example 3

Preparation of a microcrystalline plant sterol

20 kg of sitosterol (Kaukas Oy) was added as such to 180 kg of sucrose and the mixture was jet-pulverized with a dry tumbling grinder using a classifier (Oy Finnpulva Ab, grinder FP3P, classifier FPC15R). The carrier gas was air having a temperature of 63°C and a feed pressure of 2.4 bar.

Example 4

Preparation of a microcrystalline plant sterol

4530 g of sitostanol was jet-pulverized as such with a dry tumbling grinder using a classifier (Micropulva). Air was used as the carrier gas. The about 500 g coarse fraction obtained at the first stage was re-supplied to the grinder and was vibrated, whereby a very particulate product was obtained.

Example 5

Preparation of a microcrystalline plant sterol

15

Example 4 was repeated using 4100 g of sitostanol. At the first stage, 1250 g course fraction was obtained and subsequently re-supplied to pulverization, simultaneously increasing the feed rate and flushing air.

5

Example 6**Preparation of a homogenized microcrystalline plant sterol**

2000 g of microcrystalline powder prepared according to example 3 and containing 200 g of sitosterol and 1800 g of sugar was mixed with 1077 g of water and 200 g of sugar, and a homogenous suspension was formed of the substances by blending for 30 min with a blender (Braun, type 4259).

10

Example 7

15

Preparation of a homogenized microcrystalline plant sterol

2000 g of microcrystalline sitosterol-sugar powder prepared according to example 3 was mixed with 260 g of Neste 77 sugar liquid to which 538 g of water had been added (Porkkalan Sokeripuhdistamo Oy), and a homogenous suspension was formed of the substances with a blender in the above-described manner.

20

Example 8**Berry-based spread**

A product of the type of a breakfast marmalade to be spread was prepared according to the following basic recipe:

25

		g
	Water	200
30	Strawberry	500
	Sweetening agent	500
	Pectin	6.1 (+ 30 g of sugar or water)
	Potassium sorbate (20%)	6.5
	Citric acid (50%)	7.8

35

The strawberries were mashed, the mashed strawberries and water were mixed, and the mixture was cooked for a few minutes. Pectin (LM) mixed in a small amount of sugar and/or hot water was added to the mixture, which was
5 heated for a few minutes. 820 g of a sitostanol-liquid-sugar mixture prepared according to example 1 was then added and the cooking was continued for a few minutes. An acidity-regulating agent and a preservative (one or both of them can be left out, if desired) were added to the
10 mixture. The mixture was cooked further until the weight was 1150 g, whereafter the product was allowed to cool to 60°C and packed for example in jars or in containers holding one serving.

In sensory evaluation, the spread was found to
15 have an excellent structure and taste. However, some of the panel members found the mouth-feel to be a bit sandy.

Example 9

Berry-based spread

A product of the type of a breakfast marmalade to
20 be spread was prepared on the basis of example 8 by using strawberry but with 700 g of a sitosterol-liquid-sugar mixture prepared according to example 7. When the marmalade was prepared, less water had to be evaporated than in the product described in example 8, wherefore also
25 the time of cooking was shorter.

In sensory evaluation, the taste and structure of the spread were found to correspond to the product described in example 6.

Example 10

Berry-based spread

A product of the type of a breakfast marmalade to
be spread was prepared on the basis of example 9 using
strawberry but increasing the dry substance content. The
target weight was 900 g, the calculated dry substance
35 content was about 63 Bx, the actual dry substance content

was 70 Bx (the plant sterol increases the Bx value). No preservative was added.

5 In sensory evaluation, the spread was found to have a different taste than the previous product. The spread did not have a sandy mouth-feel, but both the structure and the taste were considered excellent.

Examples 11 and 12

Berry-based spread

10 A product of the type of a breakfast marmalade to be spread was prepared in the manner described in example 8, but strawberries were replaced with a corresponding amount of raspberries or boysenberries, and a sitostanol-liquid-sugar mixture prepared according to example 2 was used.

15 The products were marmalades with a beautiful colour and in sensory evaluation they were found to have an excellent colour, taste, mouth-feel and structure. There was nothing powdery or sandy in these products.

Example 13

Fruit-based spread

20 A product of the type of a breakfast marmalade to be spread was prepared according to the basic recipe described in example 8, but strawberries were replaced with a corresponding amount of orange pureed with a food press, and a sitostanol-liquid-sugar mixture prepared according to example 2 was used.

25 The pureed oranges and water were mixed and the mixture was cooked for a few mixtures. Pectin mixed in a small amount of sugar was added to the mixture and the mixture was heated for a few minutes. The stanol-sugar mixture was added and cooking was continued until the dry substance content was about 51% and the weight was about 1000 g. A preservative and an acidity-regulating agent were added to the mixture, it was allowed to cool to 80 to 35 85°C and packed for example in jars or in containers

holding one serving. The preservative and the acidity-regulating agent can be left out, if desired.

In sensory evaluation the product was found to have an excellent colour, taste, mouth-feel and structure.

5 There was nothing powdery or sandy in the product.

Examples 14 to 17

Fruit-based spread

10 A product of the type of a breakfast marmalade to be spread was prepared in the same manner as in example 13, but the oranges were replaced with a corresponding amount of apple, plum, peach or apricot pureed with a food press. The plums and the apricots were dried products that were soaked in water overnight. When apple marmalade was prepared, the amount of pectin was reduced to a half.

15 When the amount of pectin is changed, it is possible to prepare products with varying stiffnesses; due to the small amount of pectin the apple spread was therefore less thick than the other spreads. The apple spread also had a slightly pale colour; this can be
20 avoided by selecting an apple variety with a strong colour or by adding a small amount of a colouring.

Some of the panel members found the plum spread to be too solid, which can be avoided by decreasing the dry substance content of the product and/or by reducing
25 the amount of pectin.

In sensory evaluation the products were found to have an otherwise excellent taste, mouth-feel and structure. There was nothing powdery or sandy in the products. The apricot and peach spreads were considered
30 especially successful.

Example 18

Berry-based spread

A product of the type of a breakfast marmalade to be spread was prepared of strawberry in the manner
35 described in example 8, but with the mixture of a

microcrystalline plant sterol and a sweetening agent prepared according to example 3. The mixture was added as such, without forming a suspension, directly to the marmalade concoction. The mixture blended well with the concoction and no technical problems occurred.

In sensory evaluation the product was found to have an excellent colour, taste, mouth-feel and structure. There was nothing powdery or sandy in the product.

Example 19

10 Syrup

A plant-sterol-sugar mixture prepared according to example 3 was added as such to baking syrup and mixed. The pulverized mixture blended effectively with the syrup that was at room temperature and had a high dry substance content and viscosity, and the structure of the final product was very smooth. A suitable thickening agent, such as xanthan, can be added to the product, if desired, to regulate the structure to be suitable for a spread.

Example 20

20 Fat-containing emulsion

To form an emulsion, a sucrose-sitosterol mixture 90:10, ground according to example 3, was dissolved in rape-seed oil (Menu, Raisio Oy). An extra fine icing sugar correspondingly dissolved in rape-seed oil was used as a control.

The amounts of dry substance shown in Table 1 were measured into a blender. The mixture was mixed at full speed for 30 min with the blender. A homogenous emulsion was formed. The emulsions were poured into a glass container and allowed to rest at room temperature.

Table 1. Mixtures and their weighings

	Experiment	Dry substance, g	Cooking oil, g
1	sucrose-sitosterol (90:10):oil 25:75	73.19	219.48
2	sucrose-sitosterol (90:10):oil 50:50	188.96	188.62
3	icing sugar:oil 25:75	95.35	285.84
4	icing sugar:oil 50:50	216.70	216.83

Immediately after a mixing in the blender the mixtures were homogenous and milky, excluding the mixture of icing sugar and oil in the ratio of 50:50, which was a very thick mayonnaise-like homogenous mixture. The thicknesses of the mixtures depended on the proportions of oil and dry substance. Table 2 shows sensory evaluation at different times after the mixing.

Table 2. Sensory evaluation of sucrose-sitosterol-oil mixtures

	Experiment	Time			
		3 h	1 day	2 days	5 days
1	sucrose 90%: oil 25:75	homogenous, milky	homogenous, milky	separate layer of oil	separate layer of oil
2	sucrose 90%: oil 50:50	homogenous, milky	homogenous, milky	homogenous, milky	homogenous, milky
3	icing sugar:oil 25:75	homogenous, milky	separate layer of oil	separate layer of oil	separate layer of oil
4	icing sugar:oil 50:50	thick "mayonnaise"	thick "mayonnaise"	thick "mayonnaise"	thick "mayonnaise"

Example 21**Liquid margarine**

100 g of jet-pulverized (Finnpulva) sugar-plant-sterol mixture was added to 200 g of liquid margarine (Sunnuntai[®], Raisio Oy) and mixed. The mixture of sugar and plant sterol mixed well with the liquid margarine. The microcrystalline plant sterol does not form great lumps, but there is a slight mouth-feel and some small lumps. 200 g of the aforementioned liquid margarine of Raisio was added; when the dry substance content decreased, the mouth-feel became considerably smoother.

Example 22**Soft margarine**

5% of sitostanol and sitostanol-sterol preparations ground in pure form by a jet-pulverizing method according to examples 4 and 5 were mixed with a soft margarine (Keiju, Raisio). The products mixed well, forming a smooth spread, from which the powder added cannot be distinguished by the naked eye. When the spread was used, no particles were detected in the product in sensory evaluation when the product was tasted as such.

Example 23**Fat-containing emulsion**

5% of plant sterol preparations ground according to examples 4 and 5 were mixed with rape-seed oil (Kultasula, Raisio). Mixing yielded a homogenous emulsion.

Claims

1. A product containing a plant sterol, characterized by containing a microcrystalline plant sterol, a sweetening agent and water, and by being produced without admixtures.

2. A product according to claim 1, characterized in that the volumetric mean particle size of the microcrystalline plant sterol is less than 35 μm , preferably less than 30 μm and more preferably less than 25 μm .

3. A product according to claim 2, characterized in that the proportion of particles having a volumetric mean particle size exceeding 60 μm is substantially less than 10% in the microcrystalline plant sterol.

4. A product according to any one of claims 1 to 3, characterized in that the plant sterol is β -sitosterol, β -sitostanol, campestenol, clionastanol or a mixture thereof.

5. A product according to claim 4, characterized in that the plant sterol is preferably β -sitosterol or β -sitostanol, more preferably β -sitostanol.

6. A product according to any one of claims 1 to 5, characterized in that the sweetening agent is a sugar, an aqueous solution of a sugar, a starch syrup, a sugar alcohol, polydextrose or an intense sweetener.

7. A product according to claim 6, characterized in that the sweetening agent is an aqueous solution of a sugar or a sugar syrup with a dry substance content of at least about 65%.

8. A product according to any one of claims 1 to 7, characterized by being produced by pulverization and homogenization.

9. A product according to any one of claims 1 to 8, characterized by being produced using a great shear force.

5 10. A method for producing a product containing a plant sterol, characterized in that from a microcrystalline plant sterol and a sweetening agent is formed a homogenous suspension in an aqueous solution without any admixtures.

10 11. A method according to claim 10, characterized in that the crystalline plant sterol and the sweetening agent are subjected, in either order, to pulverization and homogenization.

15 12. A method according to claim 10 or 11, characterized in that the pulverization is carried out as dry grinding or wet grinding.

13. A method according to claim 12, characterized in that the pulverization is carried out as dry grinding.

20 14. A method according to claim 13, characterized in that the pulverization is carried out as dry grinding before the homogenization.

15. A method according to claim 12, characterized in that the pulverization is carried out as wet grinding.

25 16. A method according to claim 15, characterized in that the pulverization is carried out as wet grinding after the homogenization.

30 17. A method according to any one of claims 10 to 16, characterized in that the plant sterol is ground to a volumetric mean particle size of less than 35 μm , preferably less than 30 μm and more preferably less than 25 μm , for example less than 20 to 23 μm .

35 18. A method according to any one of claims 10 to 17, characterized in that the homogenization is carried out by mixing with a blender.

19. A method according to claim 18, characterized in that the homogenization is carried out by using a great shear force.

5 20. A method according to any one of claims 10 to 19, characterized in that the plant sterol is β -sitosterol, β -sitostanol, campestenol, clionastanol or a mixture thereof.

10 21. A method according to any one of claims 10 to 20, characterized in that the sweetening agent is a sugar, an aqueous solution of a sugar, a starch syrup, a sugar alcohol, polydextrose or an intense sweetener.

15 22. A method according to claim 21, characterized in that the sweetening agent is an aqueous solution of a sugar or a sugar syrup with a dry substance content of at least about 65%.

20 23. A plant-sterol-containing product, characterized by containing a microcrystalline plant sterol, fat and optionally a sweetening agent, and by being produced without any admixtures.

25 24. A product according to claim 23, characterized in that the volumetric mean particle size of the microcrystalline plant sterol is less than 35 μm , preferably less than 30 μm and more preferably less than 25 μm .

30 25. A product according to claim 23, characterized in that the proportion of particles having a volumetric mean particle size exceeding 60 μm is substantially less than 10% in the microcrystalline plant sterol.

26. A product according to any one of claims 23 to 25, characterized in that the plant sterol is β -sitosterol, β -sitostanol, campestenol, clionastanol or a mixture thereof.

27. A product according to claim 26, c h a r -
a c t e r i z e d in that the plant sterol is preferably
 β -sitosterol or β -sitostanol, more preferably β -
sitostanol.

5 28. A product according to any one of claims 23
to 27, c h a r a c t e r i z e d in that it also contains
a sweetening agent.

10 29. A product according to claim 28, c h a r -
a c t e r i z e d in that the sweetening agent is a
sugar, an aqueous solution of a sugar, a starch syrup, a
sugar alcohol, polydextrose or an intense sweetener,
preferably an aqueous solution of a sugar or a sugar syrup
with a dry substance content of at least about 65%.

15 30. A product according to any one of claims 23
to 29, c h a r a c t e r i z e d by being produced by
pulverization and homogenization.

31. Use of a product according to any one of
claims 1 to 9 or 23 to 30 as a cholesterol-lowering agent.

20 32. Use of a product according to any one of
claims 1 to 9 or 23 to 30 for the manufacture of edible
products.

25 33. Use according to claim 32 for the manufacture
of spreads, sweets, chocolate, nectars and other
beverages, yoghurts, puddings, creams, bakery products and
fats.

34. Edible products, c h a r a c t e r i z e d
in that they contain a product according to any one of
claims 1 to 9 or 23 to 30.

30 35. A spread, c h a r a c t e r i z e d in that
it contains a microcrystalline plant sterol, a sweetening
agent and berry and/or fruit.

36. A spread according to claim 35, c h a r a c -
t e r i z e d in that it also contains pectin.

Fig. 1





Fig. 2

2/2



INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 97/00585

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: A61K 9/00, A61K 31/575

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 934686 A (C.F. BOEHRINGER & SOEHNE), 21 August 1963 (21.08.63)	1-31
Y	---	32-36
A	US 5244887 A (CARL D. STRAUB), 14 Sept 1993 (14.09.93)	1-31
Y	---	32-36
A	US 4195084 A (JOHN T.H. ONG), 25 March 1980 (25.03.80)	1-36

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"B" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

17 December 1997

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Date of mailing of the international search report

09-01-1998

Authorized officer

Eva Johansson
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 97/00585

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 3881005 A (ARVIND L. THAKKAR ET AL), 29 April 1975 (29.04.75) --	1-36
A	US 4124607 A (JOHN M. BEATON), 7 November 1978 (07.11.78) --	1-36
A	WO 9219640 A1 (RAISON MARGARIINI OY), 12 November 1992 (12.11.92) -- -----	1-36

INTERNATIONAL SEARCH REPORT

Information on patent family members

02/12/97

International application No.

PCT/FI 97/00585

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 934686 A	21/08/63	BR 6237169 D	00/00/00
		DE 1131845 B	00/00/00
		DE 1139236 B	00/00/00
		LU 41386 A	15/05/62
		NL 142069 B	15/05/74
		NL 276700 A	00/00/00
US 5244887 A	14/09/93	NONE	
US 4195084 A	25/03/80	NONE	
US 3881005 A	29/04/75	AR 202034 A	09/05/75
		AU 7155674 A	29/01/76
		BE 818729 A	12/02/75
		DD 112899 A	12/05/75
		DE 2437845 A	27/02/75
		DK 427674 A	21/04/75
		FR 2240717 A,B	14/03/75
		JP 50048115 A	30/04/75
		NL 7410862 A	17/02/75
		SE 7410331 A	14/02/75
		ZA 7404587 A	25/02/76
US 4124607 A	07/11/78	NONE	
WO 9219640 A1	12/11/92	AU 664827 B	07/12/95
		CA 2102112 A	04/11/92
		DE 69127207 D	00/00/00
		EP 0594612 A,B	04/05/94
		SE 0594612 T3	
		FI 98730 B,C	30/04/97
		FI 934869 D	00/00/00
		FI 964951 A	11/12/96
		JP 6506909 T	04/08/94
		NO 933966 A	02/11/93
		PL 166991 B	31/07/95
		US 5502045 A	26/03/96